

Responses to referee 2: (received: 28 Aug 2019)

We thank the referee for very insightful questions and comments. They have helped to improve the quality of the paper. We have substantially revised and reorganized the manuscript, in many parts extended sentences and paragraphs has added. Our responses are given point-by-point below (blue Times New Roman font) following each of the reviewers' comments, which are repeated in full (black Times New Roman Italic font). Reproduced text of the revised manuscript is set in green Times New Roman font.

In addition to the change made in the manuscript to take into account your comments or the comments of the other referees, several other changes have been made and are listed here. The page number and lines indicated on the lists are taken from commenting manuscript.

Overview:

Yirdaw berthed, et al., have submitted a manuscript comparing ground-based MIR-FTS measurements of atmospheric CH₄ and N₂O at Addis Ababa, Ethiopia to that of three satellite (MIPAS, MLS and AIRS) data products. The manuscript details the Addis Ababa site, the measurements made and the spectral processing procedure (including retrieval uncertainty estimates). A brief overview of the satellite data products used are given, then coincident comparison criteria and lastly analysis of the profile and partial column comparison results.

The novelty of this manuscript is that this is the first time Addis Ababa FTIR N₂O and CH₄ measurements are compared to satellite measurements.

The manuscript content is in the scope of the AMT journal. This research will be a welcome addition to already published literature concerning FTIR data from the Addis Ababa station, and also in the wider context of atmospheric ground-based trace gases measurements (including in situ) situated on the African continent (a data sparse region of the globe). Unfortunately, the manuscript is let down in multiple critical areas and I do not recommend publication until the issues listed below are addressed; either fixed or with a sufficient logical rebuttal.

Response: We would like to thank the reviewer for this positive evaluation and critical comments that would help us to make the paper more vital to the scientific community. Furthermore, the response of each major comments and specific comments will place following each comment. I hope, all the responses given in the manuscript have satisfied the referee.

Specific comments:

S1/ AMT English guidelines and house standards: A draw-back of the submitted manuscript is that I do not believe the grammar meets the standard required for publication in AMT. The authors are referred to AMT guidelines: https://www.atmospheric-measurement-techniques.net/for_authors/manuscript_preparation.html. There are instances of incorrect grammar use, ambiguous statements (most likely a consequence of improper grammar) and repetition of statements. All such instances need to be corrected. This is no reflection on the quality of the science presented and doesn't detract (only distracts and introduces ambiguity) from the novelty and importance of the presented subject matter (along with the effort the authors have already put into the manuscript). I would have expected the more experienced co-authors to have alerted the lead author to many of these grammatical and stylistic errors. For the manuscript review, correction of such grammatical errors will be left out (to speed up the review), and only commented upon if scientific clarity is required.

Response on S1: We appreciated the referee for taking out all the concerns on the manuscript. Hopefully, all the issues raised above will be addressed in the revised manuscript.

The grammar errors have been corrected in the revised manuscript.

S2/ Could the authors clarify in the focus of the research. Comparisons are made between three satellite datasets and that of the ground-based FTIR measurements at Addis Ababa, but why? What is the motivation? In section 2.1 the manuscript alludes to why measurements at Addis Ababa are made, but only very broadly in a generic tropical atmosphere context. I gather the motivation is to use satellite measurements to validate the ground base measurements? This is unusual (usually the other way around), but a valid approach to help assess the quality of the ground-based measurements, if there is concern.

The authors state that the comparisons at the “Addis Ababa station is good to study tropical atmospheric processes” (Pg 19, L12). ‘Good’ in what context? Given the comparison results, will the ground-based CH₄ and N₂O measurements capture seasonal cycles and multi-year trends? Will biomass burning or other episodic events most likely be seen, and from what part of the tropics (the tropic is a large place)?

Response on S2: Since this result is the first to the Addis Ababa site, it required validation to assess the quality of the ground-based FTIR measurements. The motivation is to fill the gap in understanding the atmosphere over Addis Ababa, tropical region of the globe. Moreover, there are scores of measurements in tropical. The expression below has been added in section 2.1 to elaborate more about the measurement site.

Ethiopian is characterized by high rainfall and temperature variability on both spatial and temporal scales. The variability in distribution is related to altitude, latitude, humidity and winds, which are the significant factors in affecting the weather system of the country.

The sentence at pg 19, L1, “Therefore, the performance of instruments, FTIR, MIPAS and MLS in capturing CH₄ and N₂O values at the Addis Ababa station is good to study tropical atmospheric constituents.” has been rewritten as follows. Since, the paper concerns on the FTIR measurement.

Therefore, the performance of the ground based FTIR instruments in Addis Ababa, Ethiopia site would be vital to fill the scarcity of ground based FTIR measurements at tropics as the bias obtained are in agreement to other sites.

S3/ Pg 3, L19. As, in S2, the authors give a generic/broad scale reason for the importance of trace gas measurements in the tropics. I recommend that a more specific reason/motivation for Addis Ababa measurements be stated in the context of physical (or chemical) processes (emissions) related more specifically to Addis Ababa and the atmospheric footprint it ‘sees’.

Response on S3/pg3, L19: As the number of the population has increased dramatically and there is no encouraging methods for waste disposal at the Addis Abba. The following expression has been added in the manuscript as a description of the site.

Thus, the observed variation in the measurement of atmospheric trace gases would help us to understand the effects of tropical dynamics on the site. Besides, it filled the gap to the scarcity of ground based measurements in tropical.

S4/ Pg 4, L3-24. Retrieval information is incomplete, see comment T31 below. After this sentence the authors start describing the retrieval specifications (spectral Microwindows and model atmosphere layer scheme), then return to describing the optimal estimation method (L8-L24). It would be better to complete describing the retrieval theory prior to specific retrieval strategies. The information supplied in L8-L24 is ubiquitous and generic, I do not think it needs description. This section could be condensed to a single sentence stating Roger OEM approach is used (referenced) with Tikhonov regularization (reference).

Response on S4/page 4, L3-24: I think there are redundancy of concepts and we have deleted the sentences in L3-L9.

S5/ Pg4, L28: Apriori is mentioned. Are the apriori profiles used static? i.e. unvarying, or are they changing seasonally, yearly, or daily? If the apriori is static, then how is it constructed, a mean over XX years? Is the apriori based on a certain global region?

Response on S5/pg 4, L28: It was static, mean of 40 years WACCM for tropics was used.

S6/ Pg 5, Fig 1. ‘Tropics’ is a big area with a variable atmospheric state. Do the authors mean the apriori over the Addis Ababa region? Could the date of the Apriori temperature profile be put in the figure caption? Also, to show the reader the variability of the atmospheric state, could the 1-sigma SD at each layer be plotted. The authors could also possibly omit figure 1 completely, as information content is minimal.

Response on S6/pg 5, Fig 1: We agree and removed it. Similarly, the text has been changed as follows.

Both methane (CH₄) and nitrous oxide (N₂O) are well-mixed in the troposphere and their VMR decrease with height and becomes negligible with no variation above 55 km. The vertical variability of N₂O and CH₄ in the lower stratosphere is characterized by a large vertical gradient

S7/ Pg 5, L15. The description “with positive and negative signs...” can be removed. This is implicit in the retrieval. The authors should also describe the residuals. Are they dominated by random or systematic uncertainties? For instance, in Fig 2, the CH₄ fit residuals are dominated by systematic spectral error, most likely due to imperfections in the spectroscopic database line parameters.

Response on S7/pg 5, L15: The sentences “The magnitude of residuals of spectral fits are less than 1% with both positive and negative signs (CH₄: 0.4%; N₂O: 0.34%).” has been replaced by “The magnitude of the residuals of spectral fits span a range of a maximum of +0.4 % to -0.4 % for CH₄ and + 0.34 % to -0.34 % for N₂O.”

As clearly seen in the error analysis section of the manuscript pg 9 L9-21, the N₂O fit residuals are dominated by statistical error.

S8/ Pg 5, L15. The authors mention an “optimised retrieval strategy” but only give a passing mention to the Tikhonov retrieval regularization scheme. This is an important part of the retrieval; influencing overall information content and interlayer correlations of information content. Could the author please describe the Tikhonov regularization parameters. Why was the Tikhonov scheme implemented instead of using apriori uncertainties? What type of smoothing constraint is used (L1, L2 etc..), were the smoothing constraints normalised using layer thickness? what is the alpha parameter used? and how was the alpha parameter selected? is the alpha parameter static? or varies per retrieval?

Response On s8/page 5, L15: detailed description of retrieval strategy has been added after L10.

Methane and nitrous oxide vertical profiles over Addis Ababa have been obtained by fitting five and four Micro windows respectively. The retrieved state vector contains the retrieved volume mixing ratios of the target gas defined in 41 layers of the tropical atmospheric conditions. The retrieved profiles were derived using a Tikhonov-Phillips method on a logarithmic scale.

S9/ Pg 7, L4. Since equation 1 may be eliminated in section 2.2, insert equation 1 here or a reference to this equation in Rodgers, 2000.

Response on S9/pg 7, L4: The retrieved profile is a combination of a priori profile x_a and the differences of true Values and a priori weighted by the averaging kernel matrix. The mathematical expression of the retrieved profile x is given as follows.

S10/ Pg 9, L21. Section 3.3 should end and new section 3.4 “Time series” (or something similarly named) should start. The content from L22 onwards (to end of the section) is

concerned with the time series, not explicitly error estimation. The sentence starting “Concentrations of CH₄...” (Pg 6, L1) should be moved into this new section.

Response on S10/pg 9, L21: The following subsection has been added and it starts with a statement from pg 6, L1

“3.4. Time series, partial column.”

“Concentrations of CH₄ and N₂O were derived from 166 spectra of NDACC filter 3 recorded from Dec. 2009 to March, 2013.”....

S11/ Figure 6 and Pg 6, L1. Please state the reason why is only data from 2009 to 2013 is analysed? I assume the Addis Ababa station is still currently (up to 2019) taking measurements?

Response on S11/Figure 6 and page 6, L1: It has not worked since 2015 and we have taken the measurements for that period of time.

S12/ Section 4: this section details MIPAS, MLS and AIRS satellite-based measurement platforms. It would be more helpful if the focus of this section was on details about Addis Ababa overpasses for each platform (such as the number of ‘good’ overpasses as a proportion of total). This would also help diagnose if Addis Ababa is a ‘good’ site (as the authors have stated) for such satellite validation.

Response on S12/ Section 4: The time when the satellites cross Addis Ababa has to be stated under each subsection. This May use to decide the importance of the site.

S13/ Pg 11, L1. Why was only the period Mar2009 to Dec2010 used in MIPAS Addis Ababa comparisons? Why not longer?

Response on S13/pg 11, L1: This work has prepared before three years that is why we used those limited time periods. The following statement has also been added on P10, L8 to explain the time period used to validate FTIR with MLS.

The comparison of FTIR with MLS for a time period of May 2009 to February 2013 has also made. Here, in this paper we did not put anything about spatial-temporal variability of both CH₄ and N₂O

S14/ Pg 12, L3. The last two sentences, starting with “Nitrous oxide derived...” should be omitted as it refers to MLS data version 2.2, not 3.3, unless the authors state (after verifying) that the precision of MLS N₂O v3.3 is the same as v2.2.

Response on S14/pg 12, L3: The last two sentences have been written as follows:

MLS N₂O v2.2 has been validated and its precision and accuracy is respectively in Lambert et.al. (2007). The authors reported that MLS N₂O precision is 24-14 ppbv (9-41%) and the accuracy is 70-3 ppbv (9-25%) in the pressure range 100-4.6 hPa.

S15/ Section 5 details and quantifies comparisons between the satellite data products and the Addis Ababa ground based FTIR data, but (in my opinion) does not elaborate on the results with respect to other ground based FTIR site measurements. Are the biases and spread seen at Addis Ababa like that of other ground-based FTIR sites (most likely also part of the NDACC)? This would help ascertain if the Addis Ababa is a ‘good’ validation site and is network comparable. All that is required is a literature review, this will help put the results derived in this study in context.

Response on S15/ Section 5: We were set the statement due to the results of the validation and to make it clearer, it has been replaced by “Therefore, the performance of the FTIR instrument in capturing CH₄ and N₂O values at the Addis Ababa station is vital to monitor and understand the atmosphere over Addis Ababa. In addition the ground based observation has been used to supplement the satellite observations.”

S16/ Equations 4 to 10 all pertain to statistical calculations between the FTIR and MIPAS measurements. I assume the same statistical methods are applied to comparisons with the other satellite data? Maybe make this section more generic, not just MIPAS specific.

Response on S16/ Equation 4 to 10: The paragraph and equations have been rewritten as follows; sat_i (z) has been changed to X_s(z) in equation 4.

The ground based FTIR measurements of CH₄ and N₂O has been validated at different locations (e.g. Senten et al. 2008). MIPAS, MLS and AIRS have a better vertical resolution than ground-based FTIR profiles and high temporal and spatial coverage in the tropics. The analysis of the comparison between volume mixing ratio values derived from FTIR and MIPAS were performed for the data sets between March 2009 to December 2010. Furthermore, The comparison of FTIR (CH₄, N₂O) with a MLS (CH₄, N₂O) and AIRS (CH₄) for the time period of May 2009 to February 2013 has also made. Hence, the profiles from MIPAS, MLS and AIRS have been degraded to make a comparison between the FTIR and satellite observations. Therefore, the satellite measurement profiles are smoothed using the FTIR is averaging kernels of individual species obtained from the ground based FTIR retrieval by applying the procedures reported in Rodgers and Connor (2003 and given as

$$X_s = X_a + A(X_i + X_a)$$

The absolute difference at each altitude layers of a pair profile is calculated using

$$\delta_i(z) = [FTIR_i(z) - X_s(z)]$$

Where $FTIR_i(z)$ and $X_s(z)$ are the FTIR and smoothed satellite profiles of CH₄ or N₂O respectively. The mean squares error can be expressed as

$$MSE_i = \sqrt{\frac{1}{N(z) - 1} \sum_{i=1}^{N(z)} \delta_i(z)^2}$$

$$\Delta_{\text{rel}}(z) = \frac{1}{N(z)} \sum_{i=1}^{N(z)} \delta_i(z)$$

Where $\delta_i(z)$ is the difference (absolute or relative), $N(z)$ is the number of coincidences at z , $\text{FTIR}_i(z)$ is the FTIR VMR at z and the corresponding $X_s(z)$ smoothed volume mixing ratio derived from satellite instruments.

S17/ Pg 14, L3. “Hence we will focus on the random uncertainties associated with...”. This statement does not connect with the analysis in section 5.2. In section 5.2 dataset biases are quantified, which includes both random and systematic uncertainties (not separated). The standard deviations of the dataset comparisons will also include any systematic uncertainties. Maybe this sentence be retracted or changed to explain what is meant in a clearer manner.

Response on S17/pg 14, L3: To make this clear, we rewrite as follows and took only the random uncertainty of the instruments to evaluate the uncertainty of the comparison.

Here in this paper coincidence and smoothing errors are not taken into account in the full error analysis of the comparisons between remotely sensed data sets (von Clarmann, 2006). Hence, we focus on the random uncertainties of each instrument (Combined random error) that has been used to evaluate the uncertainty of the comparison (standard deviation of the difference).

S18/ Pg 14, L4. “However, the residual coincidence and horizontal smoothing errors...”. If they are important why are they not investigated? The sentence starting on L19, pg14 (“In addition, the overestimation”) also alludes (and offers conjecture) to issues arising around differences in the datasets relating to coincidence criteria but is not quantified. The authors could easily check this by changing the coincidence criteria (spatial and temporal) and see the effect of this in the dataset statistical differences.

Response on S18/pg14, L4: As the referee has suggested, the residual coincidence can be shown and we try to include on the revised manuscript.

S19/ Section 5.2 and Section 5.3. In both these sections there is no mention of how dataset degrees of freedom affect the profile differences. For example, the ground-based retrievals of CH4 have approx. 2 DOFs. Differences at different altitudes will not be independent pieces of information. At Pg 14, L25 the authors state the bias of FTIR and MLS CH4 at 18-20km is insignificant, at 17km -1.7%, and between 20-27km below 11%. Are these pieces of information independent? The authors may wish to comment on this fact and its implications.

Response on S19/ Section 5.2 and Section 5.3: No, this information is dependent

S20/ Figures 10 and 11. There is no commentary on the large ‘RD’ differences above 30km (no sensitivity?). Could the authors comment on this?

Response on S20/Figures 10 and 11: As the sensitivity of the FTIR measurement is below 27 km that indicates the retrieved value above 27 km might be from the a priori. This makes the RD large above 30km.

S21/ Pg 17, L1. *There needs to be a new section “section 5.4: Comparisons of partial columns” (or similar) starting at pg17 L1 if the authors are to start discussing partial column comparisons. Currently, the partial column comparisons for both N2O and CH4 are under section 5.3.*

Response on S21/pg 17, L1: We have added the section 5.4 as suggested by the referee.

S22/ Pg 17, L1. *Why are only MIPAS partial column comparisons conducted? What about other satellite data products?*

Response on S22/page 17, L1: We have only taken the MIPAS partial column comparison due to time limitation. However, this may consider during submitting the revised manuscript.

S23/ Fig 12. *Uncertainty/error bars could be added to all data points. This would help in assessing the comparisons.*

Response on S23/Fig 12: The middle panel of all the comparison figures have a standard error of the difference (SEMAD, blue dotted).

S24/ Pg 19, L12. *Define ‘good’, do the authors mean the measurement quality and retrievals are ‘good’, or the location, or both? Since the focus of the manuscript is on assessing the performance of the Addis Ababa FTIR measurements, explaining ‘good’ is quite important.*

Response on S24/page 19, L12: The sentence on page 19, L1, “Therefore, the performance of instruments, FTIR, MIPAS and MLS in capturing CH₄ and N₂O values at the Addis Ababa station is good to study tropical atmospheric constituents.” has been rewritten as follows. Since, the paper concerns on the FTIR measurement.

Therefore, the performance of the ground based FTIR instruments in Addis Ababa, Ethiopia site would be vital to fill the scarcity of ground based FTIR measurements at tropics as the bias obtained are in agreement to other sites.

Technical comments (no particular order):

T1/ Title: *FTIR should be expanded, not an acronym. There is no need for the chemical formulas. The word ‘measurements’ should also be added after FTIR. So...I recommend the full title should be along the lines of: “Methane and nitrous oxide from ground-based Fourier transform infrared spectrometer measurements at Addis Ababa: observations, error analysis and comparison with satellite data.”*

Response T1, title: The title has been changed to Methane and nitrous oxide from ground-based Fourier Transform Infrared spectroscopy measurements at Addis Ababa: observations, error analysis and comparison with satellite data.

T2/ Pg 1, L2. Possible change: “total column abundances and vertical distribution of various constituents in the atmosphere” to “total column trace gas abundances and vertical distributions”.

Response on T2, P1, and L2: have been replaced by “total column trace gas abundances and vertical distribution.”

T3/ Pg 1, L4. The superlative sentence “They reveal the high quality of FTIR measurements at Addis Ababa” is not required. The data and analysis reveal this.

Response on T3, P1, L4: has been changed to “The data and analysis reveal the high quality of the FTIR measurements.”

T4/ In the abstract, I do not think it is necessary to specify satellite data product versions, for example ‘V5R_CH4_224’. This is done in the main body.

Response on T4: The importance of describing the version of the data on the abstract only to make it clear the readers with what version of the data has been validated. However, if the reviewer does not agree, we change it on the revised manuscript.

T5/ Pg 1, L12. There are phrases throughout the manuscript of the sort “a positive bias of less than 0.14 ppmv (9%) is found in the altitude range of 21to 27 km”. I gather this means there is a maximum positive bias of 0.14 ppmv in the range 21 to 27km? This may be a better way to state it.

T6/ Pg 2, L1. CH₄, N₂O and CFCs are also stratospheric species...;)

Response on T6, P2, L1: has been rewritten as follows

"Methane (CH₄), nitrous oxide (N₂O) and chlorofluorocarbons (CFCs) are the main source gases to the chemical families NO_x, ClO_x, and HO_x at troposphere and stratosphere (Jacobson, 2005)."

T7/ Pg 2, L7. “ENIVSAT” to “ENVISAT satellite”.

Response T7, P2, L7: has been added a word following "ENVISAT" "satellite"

T8/ Pg 2, L9. Remove the word ‘recent’ from “The recent increasing...” (also replace ‘to the’ with ‘on’).

Response T8, P2, L9:Done

T9/ Pg 2, L10. Merge the sentences to read: “The recent increasing impact of CH₄and N₂O to the global warming has also been assessed by the last AR4 IPCC report (IPCC, 2007; Sussmann et al., 2012), additionally N₂O will become the dominant ozone depleting substance emitted in the 21st century (Ravishankara et al., 2009).”

Response T7, P2, L9-11: the sentence has been changed to

The increasing impact of CH₄ and N₂O on global warming has also been assessed by the last AR4 IPCC report (IPCC, 2007; Sussmann et.al. 2012), Additionally N₂O will become the dominant ozone depleting substance emitted in the 21st century (Ravishankara et.al. 2009).

T10/ Pg 2, L11. What is IASI? Expand to say: "IASI instrument aboard the MetOp-2 satellite". MetOp-1 or MetOp-2...I can't remember.

Response T10, P2, L11: The sentence has been started with the following

"The Infrared Atmospheric Sounding Interferometer (IASI) on-board METOP-1"

T11/ Pg 2, L18. Rephrase first sentence: "In the tropics two important..."

Response T11, P2, L18: The sentence has been rephrased as follows.

Two important exchange processes are taking place in the tropics, those are the inter hemispheric exchange and entry of tropospheric air mass to the stratosphere (Petersen et.al. 2010; Fueglistaler et.al. 2009).

T12/ Pg 2, L25. Replace 'launched' with 'taken'.

Response to T12, P2, L25: "launched" has been replaced by "taken"

T13/ Pg 2, L27. Replace "The quality of ground based FTIR measurements" with "The Addis Ababa FTIR measurements".

Response on T13, P2, L27: has rewritten as follows

The Addis Ababa FTIR measurement of atmospheric trace gases and their use to understand various lower and middle atmospheric processes have been reported in a number of previous studies (Takele Kenea et.al., 2013; Mengistu Tsidu et.al., 2015; Schneider et.al., 2015, 2016; Barthlott et.al.,2017).

T14/ Pg 2, L33. Replace 'confirm' with 'show'.

Response in T14, P2, and L33: "confirmed" has been replaced by "showed"

T15/ Pg 2, L34. Replace 'biased high and provided +14% as the most likely bias' with 'biased 14% high'.

Response on T15, P2, L34: The sentence has rewritten as follows

Laeng et.al. (2015) found the MIPAS CH₄ profiles V5R_CH4_222 below 20 to 25 km biased +14% high

T16/ Pg 2, L33. The reference Kenea throughout the manuscript should be replaced with Takele Kenea (2013)?

Response: "kenea" has been replaced by "takele kenea" though out.

T17/ Pg 2, L33. The quoted references of Laeng (2015) and Plieninger (2016) refer to MIPAS comparisons with other satellite products. The paragraph starting at Pg 2, L25 concerns Addis Ababa FTIR measurements. There is a jump in topic. Reading as is, it could easily be taken that Addis Ababa measurements were used in these studies. These sentences should be removed or moved to a different part of the manuscript.

Response on T17, P2-3: The last two statements on paragraph 4 of introduction have moved to the subsection 4.1, since those sentences have been briefed about the bias of MIPAS measurements.

T18/ Pg 3, L3. The sentence “In this study, the previous work on intercomparison is extended to source gases CH₄ and N₂O from ground-based FTIR” is quite ambiguous. Either remove or make more specific to Addis Ababa.

Response on T18, P3, L3: To make it clear, the statement has been rephrased as follows

In this study, the previous work on intercomparison of ozone (Takele Kenea et al., 2013) and water vapour. ((Samuel Takele Kenea., 2014)) are extended to source gases CH₄ and N₂O from ground-based FTIR of Addis Ababa site.

T19/ Pg 3, L7. “approach” can be replaced with ‘strategy’.

Response on T19, P3, L7: The word "approach" has changed to "strategy"

T20/ Pg 3, L13. Is Addis Ababa part of the Network for the Detection of Atmospheric Composition Change (NDACC)? I suspect so, if this is the case it should be stated. The Takele-Kenea paper should be used as a site reference paper.

Response on T20, P3, L13: Our FTIR observatory has not registered yet as part of NDACC. Moreover, the reference has been added as suggested by the referee.

T21/ Pg 3, L15. Could the Addis Ababa site altitude (MASL) also be added?

Response on T21, P3, L13: Done

T22/ Pg 3, L15. How is ‘suitability’ defined? Why is it suitable? Could possibly mention the amount of cloud free days a year.

Response on T22, P3, L15: It is known that Ethiopia is over 250 days per year is cloud free.

T23/ Pg 3, L18. The superlative “extremely” can be removed, not needed.

Response on T23, P3, L18: Done

T24/ Pg 3, L23. The superlative “very successfully” can be removed, not needed.

Response on T24, P3, L23: Done, we have been rewritten the sentence as

“Fourier transform spectroscopy has been applied successfully to study trace gases in the atmosphere by examining atmospheric absorption lines in the infrared spectrum from solar.”

T25/ Pg 3, L24. Replace 'sun' with 'solar'

Response on T25, P3, L24: Done (see the above response)

T26/ Pg 3, L27. The sentence "This technique..." should be moved to precede the sentence "The high resolution..."

Response on T26, P3, L27: We exchanged the two sentences

This technique uses the Sun as a light source to quantify molecular absorptions in the atmosphere and then retrieve trace gases abundance. The high-resolution FTIR Spectrometer, Bruker IFS120M upgraded with 125M electronics, from the Bruker Optics Company in Germany was installed in May, 2009 at the Addis Ababa site.

T27/ Pg 3, L28. "Using seven narrow band filters". Assuming Addis Ababa is an NDACC site, do the seven filters meet NDACC specifications?

Response on T27, P3, L28: Yes

T28/ Pg 3, L29. It is mentioned an InSb detector is used to take measurements in over the range: 1500-4400cm⁻¹

. There is no mention of detectors used to measure down to 750cm⁻¹, as mentioned in the prior sentence. Are measurements taken below 1500cm⁻¹?

Response on T28, P3, L29: We have removed "The spectral coverage of the IFS120M instrument at the Addis Ababa site is 750 - 4000 cm⁻¹ using seven filters." sentence from the manuscript.

T29/ Pg 3, L31. Replace "we used PROFFIT V...algorithm", with "we used the retrieval code PROFFIT (Ver95)".

Response on T29, P3, L31: Done

T30/ Pg 4, L2. As the sentence reads, PROFFIT was developed to only retrieve CH₄ and N₂O. Could this sentence be corrected to reflect the fact PROFFIT was developed to retrieve multiple species.

Response on T30, P4, L2: The phrases in green are added in the sentences below

It has been developed based on semi-empirical implementation of the Optimal estimation Method (Rodgers, 2000) to derive the VMR profiles and column amounts of multiple species. Hence, CH₄ and N₂O profiles from measured spectra in the microwindows that span spectral range of 2400 - 2800 cm⁻¹ have been discussed in this paper.

T31/ Pg 4, L3. The sentence "This algorithm..." only tells half the information. Once a forward model calculation is completed, what happens next?

Response on T31, P4, L3: "to produce the synthesized spectra" is added before we closed the statement.

T32/ Pg 4, L4. At the end of the sentence “The vertical profiles...N2O respectively” change to “N2O respectively (see table 1 for spectral regions).” and could possibly be moved to section 3.1.

Response on T32, P4, L4: This has done by taking “The vertical profiles over Addis Ababa have been obtained by fitting five and four selected spectral regions for CH₄ and N₂O respectively (see table 1 for spectral region).” to P5 L6.

T33/, Pg4, L6. Could the bottom (base) layer height be stated.

Response on T33, P4, L6: We have added “, 2.45 to 85 km” after 41 layers to show the lower and top level of the layers.

T34/ Pg4, L27. replace ‘setup’ with ‘strategy’

Response on T34, P4, L27: Done

T35/ Pg 4, L29. This is the first time the ‘NDACC’ and ‘IRWG’ acronyms are used, please state in full.

Response on T35, P4, L29: We agree that NDACC that has been stated in page 5 would be changed to page 4, L29.

T36/ Pg 5, L4. I think the sentence “The vertical variability...” is not required, or more informationis required, i.e. define ‘large’.

Response on T36, P5, L4: As the referee suggested to delete Figure 1 and to make the explanation more clear, the paragraph has been replaced by “Both methane (CH₄) and nitrous oxide (N₂O) are well-mixed in the troposphere and their VMR decrease with height and becomes negligible with no variation above 55 km. The vertical variability of N₂O and CH₄ in the lower stratosphere is characterized by somewhat higher vertical gradient as compared to the other layers.”

T37/ Pg 5, L8. “The micro-windows have been adopted from different sources”. Why, and could the sources, please be referenced.

Response on T37, P5, L8: We mean, micro windows that give us a minimum residuals and errors were considered. The references from where the micro windows are adopted has been added at P5, L8.

The microwindows have been adopted from different sources (. Senten et al. 2008; Sussmann et al., 2011; Arndt et al., 2004).

T38/ Pg, 5, L9. Modified Microwindows: why is this?

Response on T38, P5, L9: The reasons why we modify the micro windows are due to high residuals obtained between the measured and synthesized spectra at the Addis Ababa site. Thus, the micro-windows recommended by the NDACC might be useful for the other FTIR

sites found in mid and high latitudes. The references from where the micro windows are adopted has been added at P5, L8.

The microwindows have been adopted from different sources (.Senten et al. 2008; Sussmann et al., 2011; Arndt et al., 2004).

T39/ Pg 5, L7. After the end of the sentence “lines are presented”, the sentence Pg 4, L4 “The vertical profiles...” should be inserted.

Response on T39, P5, L7: The sentence has added

The vertical profiles over Addis Ababa have been obtained by fitting five and four selected spectral regions (microwindows) for CH₄ and N₂O respectively.

T40/ Pg 5, L12. Remove the word “example”. Also, could the authors detail the Signal to Noise ratio (SNR) of the Addis Ababa spectra. All this information is part of the ‘optimization process’.

Response on T40, P5, L12: Done

T41/ Table 1. could “int. Gases” be replaced with “Interfering gases”. In the table legend, could “column amounts” be replaced with “total column amounts”. Just an idea, (authors discretion), could another column be added for DOFs of the 0-27km partial column.

Response on T41, Table 1: Done

T42/ Figure 2. The usual convention is for measurement spectra data to be displayed as points (usually joined) and simulations as thin lines. The authors have the opposite of this. A minor point, and up to the authors if they would like to change it a more standard convention.

Response on T42, Figure 2: Done

T43/ Figure 2 Legend. Acronyms are not explained prior, so the full name is required. Sorry. SEA is an unusual way to label/present the solar angle. Does SEA mean Solar elevation angle? The standard convention is solar zenith angle (SZA = 90.0 - SEA). I recommend that SZA be presented, not SEA. Spectra time is presented as “101715”, please reformat to 10h17m15s or similar. Is SD the root mean square difference of the measured spectra and forward model? I.e. RMS. Please define.

Response on T43, Figure 2: Yes, the solar zenith angle is 20.6° but we expressed in terms of solar elevation angle 69.4°. All the referee comments has taken to the caption of figuer 2 and 3. The RSM is defined as a set of values is the square root of the arithmetic mean of the squares of the values,

T44/ Pg 7, L2. To be pedantic, “the most” can be replaced with “an” (as a matter of opinion).

Response on T44, P7, L2: Done

T45/ Pg 7, L11. The authors should state at this point the units of the AVKs displayed, normalised to layer VMR or not? i.e. [VMR/VMR]

Response on T45, P7, L11: Figure 4. Has been changed to the selected altitude of AVKs and its unite has shown in the figure (x-axis).

T46/ Pg 7, L16. Pedantic point, but any AVK that has non-zero elements has 'sensitivity', that is, infer information from the spectra. 0.5 is an arbitrarily defined 'cut-off'. So, I think a better statement would be "Fig 4 (top panel) shows that the.... has a sensitivity greater than 0.5 over the altitude range 2.45km to 27km".

Response on T46, P7, L16: Changed

Fig 4 shows that the retrieval of CH₄ and N₂O have a sensitivity greater than 0.5 over the altitude range 2.45km to 27km.

T47/ Pg 8, L2. Could the altitude ranges of the two independent partial columns be stated.

Response on T47, P8, L2: It is the troposphere, below 17 km and the altitude range 18-27 km.

T48/ Pg 8, L4. The sentence "The amplitude..." repeats information given earlier so can be removed.

Response on T48, P8, L4: It was not explained in the FTIR and retrieval section.

T49/ Pg 8, L5. The sentence "We also ignore": SorryI cannot understand this sentence (includes spelling mistakes), can it please be reworded and made clearer. The cited reference (Rinsland, 2005) does not contain any such information pertaining to altitude resolution. Can the authors please check the reference is correct or point out where it is in the paper.

Response on T49, P8, L4:

T50/ Figure 4 legend. The date time stamp of spectra the analysis is performed on should be given.

Response on T50, Figure 4: The figure has changed and the date has also shown in the caption.




Figure 3. Sensitivity analysis of the retrieved profiles of CH₄ (left) and N₂O (right) at Addis Ababa using the selected rows of the averaging kernels as a function of altitude. The dotted lines are the sum of the rows of the averaging kernels for a spectrum measured on Feb. 26, 2013 for CH₄ and Dec 31, 2009 for N₂O.

T51/ Pg 9, L4. Could the acronyms for sources of error be included, i.e. ‘instrument line shape (ILS)’, so that they correspond to the legend labels in fig5.

Response on T51, P9, L4: Yes, it was included in the figure and represented by a red line, but it was overlapped by other lines.

T52/ Pg 9, L5. Can the term ‘zero baselines offset’ be ‘zero level baseline offset’

Response on T52, P9, L5: yes, it is zero level baseline offset.

T53/ Pg 9, L5. I assume statistical error means ‘random’ error, maybe rephrase as “statistical (random) error”. Also take out “typical” as it relates directly to an example.

Response on T53, P9, L5: Done

T54/ Pg 9, L17. Change part of the sentence “for the profile”, to “for the N2O profile”.

Response on T54, P9, L17: Done

T55/ Pg 10, L6. The sentence starting “Vertical resolution...” needs to be referenced.

Response on T55, P10, L6: The vertical resolutions that has been stated in the discussion manuscript was for the old version. This has replaced by

The vertical resolution of MIPAS ranges from 2.5 to 7 km for CH₄, and from 2.5 to 6 km for N₂O in the reduced-resolution period (Plieninger et al., 2015).

T56/ Pg 10, L8. The sentence starting “The analysis of the comparison...” should be moved to section 5.1.

Response on T56, P10, L8: As the referee has stated, the expression has already in section 5.1 and then we have removed it from this subsection. (see the response on S16).

T57/ Pg 11, L17. The starting sentence should be reworded: “...we have used the MLS N2O (v3.3) product to validate the ground-based Addis Ababa FTIR measurements.”

Response on T57, P11, L17: The sentence in pg 11, L17 has been replaced by

“MLS N₂O data set has been used to validate the ground-based FTIR measurements. However, methane (CH₄) data are derived using coincident measurements of atmospheric water vapor (H₂O), carbon monoxide (CO) and nitrous oxide (N₂O) from the EOS MLS instrument on the NASA Aura satellite and detail are given in Minschwaner et al. (2015).”

T58/ Pg 11, L18. The altitude levels are given in pressure coordinates. So far in the manuscript, the altitude units have been in kilometres. Could the authors include the geometric altitude as well as the pressure. I.e. “100 and 0.1 hPa (XX to XX km’s)”

Response on T58, P11, L18: those altitude ranges are showing only for the data set and in the comparison we put them in km as it is known that the altitude can be expressed both in pressure and km, such as 100-0.1hpa mean 17-63 km.

T59/ Pg 12, L7. This sentence needs a reference.

Response on T59, P12, L7: The sentence has rewritten as follows and reference was also added.

Operating in nadir sounding geometry, the Atmospheric Infrared Sounder (AIRS) on board the Aqua satellite launched into Earth orbit in May 2002 Chahine et al. (2006).

T60/ Pg 12, L11. Unfortunately, the sentence starting “The spectral resolution...” does not make sense, could this be reworded, rephrased. The following sentence can be omitted as it does not add any information concerning this study.

Response on T60, P12, L11: we changed the paragraph as follows.

Operating in nadir sounding geometry, the Atmospheric Infrared Sounder (AIRS) on board the Aqua satellite launched into Earth orbit in May 2002 Chahine et al. (2006). AIRS is a medium-resolution infrared grating spectra radiometer and a diffraction grating disperses the

incoming infrared radiation into 17 linear detector arrays comprising 2378 spectral samples. The satellite crosses the equator at approximately 1:30 A.M. and 1:30 P.M. local time, resulting in near global coverage twice a day. AIRS 2378 channels covers from 649 to 1136, 1217–1613 and 2169–2674 cm^{-1} . It also measures trace gases such as O₃, CO and to some extent CO₂. AIRS CH₄ and N₂O retrievals have been characterized and validated by Xiong et al. (2008) and Xiong et al.(2014) respectively.

T61/ Pg 12, L13. *Could the data product version of AIRS CH4 be added?*

Response on T61, P12, L13: AIRS version 6

T62/ Pg 12, L20. *The word ‘version’ can be removed.*

Response on T62, P12, L20: Done

T63/ Pg 12, L22. *I think the meaning of ‘degraded’ means smoothed, so maybe the sentence could read “...MLS have been degraded (smoothed) to make a ...” or replaced degraded with smoothed.*

Response on T63, P12, L22: Done

T64/ Pg 12, L29. *Replace ‘parameters’ with ‘statistics’.*

Response on T64, P12, L29: Done

T65/ Pg 13, L11. *I assume that Sati(z) is smoothed? So maybe state that: “and the corresponding Sati(z) smoothed volume mixing ratio is derived from...”*

Response on T65, P13, L11: Equation 4 to 10: The paragraph and equations have been rewritten as follows; $\text{sat}_i(z)$ has been changed to $X_s(z)$ in equation 4.(see Response on S16).

T66/ Figures 7, 8 and 9 could be combined into a single 3x3 figure, so could figures 10 and 11 (2x2)

Response on T66, Figure 7: They would not see clearly.

T67/ Pg 14, L26. *“In the tropopause layer”, could the tropopause layer be defined, i.e. “In the tropopause layer (~XX-XX km)”*

Response on T67, P14, L26: It has been defined as 16-18 km. In addition, it varies in height according to season. Here, in this work we have already put it in the next expression 17 km.

T68/ Figures 7,8,9,10 & 11. *The legends in the figures are slightly different. Maybe standardise these as the figure captions all reference back to fig 7 caption. Figure 7 caption should also explain/define the legend captions. For example: “mean difference FTIR minus MIPAS (MAD, blue solid line) ...”*

Response on T68/ Figures 7, 8 9 10 & 11: The legends in the figures have been corrected and all the legends have also stated in the caption.

Figure 7. Comparison of CH₄ from MIPAS reduced resolution (V5R_CH4_224) and FTIR. Left panel: mean profiles of MIPAS (red) and FTIR (black) and their standard deviation (horizontal bars). Middle panel: mean difference FTIR minus MIPAS (**MAD**, blue solid), standard error of the difference (**SEMAD**, blue dotted), and mean relative differences FTIR minus MIPAS relative to their averaged (**MRD**, green, upper axis). Right panel: combined mean estimated statistical error of the difference (**combined error**, red dotted, contains MIPAS instrument noise error and FTIR random error budget), standard deviation of the difference (**STDMAD**, black solid).

T69/ Pg 16, L2. The first sentence could be abbreviated to “FTIR N₂O mixing ratio MIPAS comparison results are shown in fig 10.”

[Response on T69, P16, L2: Done](#)

T70/ Pg 16, L8. (-0.02 ppmv) ... include units please.

[Response on T70, P16, L8: Done](#)

T71/ Pg 16, L11. Replace ‘can’ with ‘could’.

[Response on T71, P16, L11: Done](#)

T72/ Pg 16, L17. Add to end of sentence: “the value derived from the FTIR is overestimated (relative to MLS)”

[Response on T72, P16, L17: Done](#)

T73/ Pg 18, L4. The last part of the first sentence “..., which is a very useful...” is not needed.

[Response on T73, P18, L4: has been corrected](#)

T74/ Figure 12. The date label lacks information on the years(s), only months are given. Add information on the year(s) comparisons were made.

[Response on T74, Figure 12: It has been corrected by adding May 2009 to December 2010.](#)

T74/ Acknowledgement: Remove full stop at the start of the first sentence. Remove the word ‘besides’. Support, not supports.

[Response on T74: Done](#)

T75/ References. The authors may wish to take out, or update references concerning an ACPD articles, such articles have not passed the peer review process. Reference formatting differs, so would be good to get it all consistent. Decide on an author convention for Samuel Takele Kenea, as this author is referenced a few times, but referenced differently (Takele Kenea S.).

[Response on T75, References: All the reference with ACPD, Atmos. Meas. Tech. Discuss. and in naming Takele Kenea has been corrected.](#)

