

## Supplementary


### S1 Composition of NPL calibration standard mixture

Table S1 : Composition of NPL calibration standard mixture (D09 0523, June 2014)

Compounds	nmol/mol	Expanded uncertainty	Compounds	nmol/mol	Expanded uncertainty
ethane	4.22	0.08	isoprene	4.19	0.09
ethene	4.06	0.08	heptane	4.18	0.08
propane	4.25	0.09	benzene	4.00	0.08
propene	4.18	0.08	2,2,4-trimethylpentane	4.15	0.08
2-methylpropane	4.37	0.11	octane	4.2	0.08
butane	4.14	0.08	toluene	3.99	0.1
ethyne	4.3	0.21	ethylbenzene	3.99	0.1
trans-but-2-ene	4.14	0.08	m-xylene	4.04	0.11
but-1-ene	4.14	0.08	p-xylene	3.98	0.1
cis-but-2-ene	4.19	0.08	o-xylene	3.99	0.1
2-methylbutane	4.21	0.08	1,3,5-trimethylbenzene	4.09	0.11
pentane	4.21	0.08	1,2,4-trimethylbenzene	4.03	0.11
1,3-butadiene	4.13	0.08	1,2,3-trimethylbenzene	4.16	0.11
trans-pent-2-ene	4.13	0.08	$\alpha$ -pinene	3.88	0.08
pent-1-ene	4.14	0.08	$\beta$ -pinene	3.95	0.2
2-methylpentane	4.23	0.08	limonene	3.99	0.12
hexane	4.15	0.08			

## S2 MnO<sub>2</sub> and thiosulfate scrubber BVOCs test results



As seen on Figure S1, the monoterpenes were unaffected by the scrubber (with a relative deviation of less than  $\pm 10\%$ ) but for oxygenated compounds, a loss of signal remained even after nine hours: e.g. nopinone (12%), citral (23%), linalool (38%) and menthol (45%).



5

**Figure S1 : Relative deviation to direct measurement of VOC concentrations when sampling with a MnO<sub>2</sub> scrubber. Each bar represents one measurement.**


### S3 Uncertainties repartition in field campaign conditions



5 **Figure S2 : Uncertainty repartition for the five terms considered (precision, calibration, integration, memory effect and scrubber) at (a) 2 ppt and (b) 100 ppt with field campaign analytical parameters.**

## S4 Concentrations time series for the 20 BVOCs measured during the LANDEX-episode 1 field campaign in summer 2017

5 BVOC time series of  $\beta$ -pinene,  $\alpha$ -pinene, limonene, myrcene,  $\Delta^3$ -carene, linalool, isoprene and  $\beta$ -caryophyllene are presented on the main publication. The other compounds are presented on Figure S2.



10 **Figure S3 : Times series of BVOC concentrations with their associated uncertainties ( $k=2$ ) for a selection of compounds observed during the LANDEX field campaign (camphene, p-cymene, nopinone, citral, sabinene, terpinolene, ocimene,  $\gamma$ -terpinene,  $\alpha$ -terpinene,  $\alpha$ -phellandrene, 2-carene).**